

Iron(II) Triflate Catalyzed Synthesis of 3,4-Dihydropyrimidin-2(1H)-ones via the Biginelli Reaction

Kyle A. Lyth, Rem Quintin V. David, Nicholas M. Edelen, Habiba Hamitova, Magnum L. McManigal, Rylan B. Nelson, Ram S. Mohan*

Laboratory for Environmentally Friendly Organic Synthesis, Department of Chemistry, Illinois Wesleyan University, Bloomington, IL, USA

Email: *rmohan@iwu.edu

How to cite this paper: Lyth, K.A., David, R.Q.V., Edelen, N.M., Hamitova, H., McManigal, M.L., Nelson, R.B. and Mohan, R.S. (2026) Iron(II) Triflate Catalyzed Synthesis of 3,4-Dihydropyrimidin-2(1H)-ones via the Biginelli Reaction. *Green and Sustainable Chemistry*, 16, 60-67.
<https://doi.org/10.4236/gsc.2026.162004>

Received: April 12, 2026

Accepted: May 23, 2026

Published: May 26, 2026


Copyright © 2026 by author(s) and Scientific Research Publishing Inc.
This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).
<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

The synthesis of dihydropyrimidinones is of interest due to their biological activities. A common method for their synthesis is the Biginelli reaction, which is a one-pot multicomponent reaction of an aldehyde, urea and a β -ketoester. The Biginelli reaction is typically catalyzed by a Brønsted or Lewis acid. However, many of these catalysts, such as $\text{BF}_3 \cdot \text{Et}_2\text{O}$ and AlCl_3 are corrosive and/or toxic. Herein, we report the iron(II) triflate-catalyzed (10.0 mol%, 22 - 24 h) synthesis of dihydropyrimidinones via the Biginelli reaction in isopropanol as a solvent. The reaction is also reported in two other solvents, benzotrifluoride (a useful replacement for CH_2Cl_2) and 2-methyltetrahydrofuran, a bio-derived solvent. Iron(II) salts are attractive catalysts because of their low cost, low toxicity, and ease of handling.



Keywords

Aldehydes, Biginelli Reaction, 3,4-Dihydropyrimidin-2(1H)-ones, Green Chemistry, Iron(II) Triflate, Multicomponent Reactions

1. Introduction

Dihydropyrimidinone scaffolds are of considerable interest in medicinal chemistry

because of their relative ease of construction and the range of biological activities they exhibit. They have been reported to show anti-hypertensive and calcium-channel blocking activities [1], anti-mitotic properties [2], antioxidant properties [3], and antibiotic properties [4]. Although Biginelli reported the first synthesis of dihydropyrimidinones in 1893 [5], it is only since the 1990's that several other methods for their synthesis have been reported. The Biginelli reaction is a classic example of a multicomponent reaction (MCR) that allows rapid assembly of a complex structure in a single step [6] [7]. The mechanism of the Biginelli reaction has been extensively studied [8] [9]. In the last couple of decades, several catalysts have been reported for the Biginelli reaction, which include Lewis acids and bases. A few representative examples of Lewis acid/acidic catalysts used for the Biginelli reaction include *tetra*-butyl ammonium bromide [10], CaF_2 [11], $\text{Y}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ [12], $\text{Bi}(\text{OTf})_3$ [13], $\text{Cu}(\text{OTf})_2$ [14], and InCl_3 [15]. The Biginelli reaction has also been done under solvent-free conditions using montmorillonite KSF [16]. There are fewer reports of the Biginelli reaction catalyzed under basic conditions, and these include $(\text{NH}_4)_2\text{CO}_3$ [17], PPh_3 [18], and *t*-BuOK [19]. Organocatalysts, such as L-proline [20], have also been used to catalyze the Biginelli reaction. A few examples of the asymmetric Biginelli reaction have been reported as well [21]. Iron compounds have attracted attention as catalysts in recent years due to their ready availability and relative non-toxicity compared to precious metal-based catalysts [22]. Iron(III) salts such as $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ [23], FeF_3 [24], $\text{Fe}(\text{CF}_3\text{CO}_2)_3$ [25], $\text{Fe}(\text{CF}_3\text{SO}_3)_3$ [25], and $\text{Fe}(\text{OTf})_3$ [26] have been successfully employed as catalysts for the Biginelli reaction. Iron(III) chloride worked well as a catalyst, but the reaction required 1.5 equivalents of urea for optimal results, and the isolation protocol generates considerable aqueous waste. Iron(III) fluoride also works well as a catalyst, but the product isolation protocol requires an aqueous work-up, thus resulting in a considerable aqueous waste stream that can contain HF. In addition, FeF_3 itself is quite toxic [27]. Iron(III) acetate and iron(III) triflate catalyzed reactions have been reported under solvent-free conditions, but again, the product isolation protocol generates a lot of aqueous waste stream. A solvent-free protocol has been reported using $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ [28]. However, caution must be exercised in grinding metal nitrates, especially on a large scale. Sustainable approaches to the Biginelli reaction have been summarized in the literature [29].

2. Results and Discussion

Due to our continued interest in developing environmentally friendly synthetic methodology, we investigated the utility of iron(II) trifluoromethanesulfonate (triflate) as a catalyst for the Biginelli reaction. Iron(II) salts are generally inexpensive, not very corrosive, and easy to handle. There are fewer reports of the utility of iron(II) triflate in organic synthesis [30]–[33]. Herein, we report the utility of iron(II) triflate as a catalyst for the Biginelli reaction that minimizes the generation of an aqueous waste stream.

The results of this study are summarized in **Table 1**. As can be seen from **Table 1**, iron(II) triflate (10.0 mol%) is an efficient catalyst for the synthesis of dihydropyrimidinones from a variety of aldehydes. Although the reaction worked with lower catalyst loading (5.0 mol%), optimal reaction times and yields were seen with the use of 10.0 mol% catalyst. Several solvents were examined for the reaction, and the best results were obtained in isopropanol, which is a relatively environmentally benign solvent [34]. This methodology avoids an aqueous work-up and hence large aqueous waste streams are avoided. The yields were lower with furfural (entry 1h) and decanal (entry 1i). The results with aldehydes containing an electron-withdrawing group, such as 3-nitrobenzaldehyde and 3-hydroxybenzaldehyde, were less promising, and the crude product was a mixture of desired product and several intermediates. We also explored the utility of an aprotic solvent, benzotrifluoride (BTF) [35] [36]. BTF is an attractive and safer alternative to the highly toxic CH_2Cl_2 , which is now banned by the US EPA. In addition, the reaction was also attempted in 2-methyltetrahydrofuran (2-MeTHF), which is a bio-derived solvent readily available from furfural and levulinic acid, and is considered an environmentally friendly alternative to THF [37]. The use of 2-MeTHF as a solvent gave better yields than BTF.





Table 1. Iron(II) triflate catalyzed synthesis of 3,4-dihydropyrimidin-2(1*H*)-ones via the Biginelli reaction.

$\text{1a-i} + \text{2} + \text{3} \xrightarrow[\text{70 } ^\circ\text{C, 22-24 h}]{\text{Fe(OTf)}_2 \text{ (10.0 mol\%)}} \text{4a-i}$

(1.0 eq) (1.1 eq) (1.1 eq)

Entry ^a	Aldehyde ^b	Solvent ^{c,d}	Yield ^e (%) 4a - i
1a [38]		ⁱ PrOH	72
		BTF	61
		2-MeTHF	74
1b [38]		ⁱ PrOH	93
		BTF	59
		2-MeTHF	74
1c [38]		ⁱ PrOH	73
1d [39]		ⁱ PrOH	85 ^f

Continued

1e [38]		ⁱ PrOH	90
		BTF	73
		2-MeTHF	86
1f [40]		ⁱ PrOH	73
1g [41]		ⁱ PrOH	73
1h [38]		ⁱ PrOH	40
1i [41]	CH ₃ (CH ₂) ₈ CHO	ⁱ PrOH	56

^aSuperscript against entry number refers to literature reference for product. ^bAll aldehydes are commercially available and were purified before use. ^cReaction mixtures were heated using a temperature-controlled hot plate. ^dThe progress of the reaction was followed by TLC (disappearance of aldehyde, 2,4-DNP stain). ^eRefers to yield of the isolated product that was determined to be $\geq 98\%$ pure by ¹H & ¹³C NMR. All products have been reported previously and were characterized by ¹H & ¹³C NMR spectroscopy. Reactions were run at least two times and yields were found to be reproducible. ^fReaction was run with 1.3 equivalents of ethyl acetoacetate and 1.3 equivalents of urea.

3. Experimental Section

Three representative procedures are given here.

Reaction in isopropanol (entry 1b):

A solution of *p*-anisaldehyde (0.212 g, 1.56 mmol, 1.0 eq), urea (0.103 g, 1.72 mmol, 1.1 eq) and ethyl acetoacetate (0.223 g, 0.218 mL, 1.72 mmol, 1.1 eq) in isopropanol (4.0 mL) was stirred at room temperature as iron(II) triflate (0.055 g, 0.156 mmol, 10.0 mol%) was added. The reaction mixture was heated at 70 °C (temperature-controlled hot plate). After 22.5 h, the mixture was cooled to room temperature. The solvent was removed on a rotary evaporator, and the residue was triturated with CH₃OH/H₂O (1:1, v/v, 5.0 mL) once to yield 0.42 g (93%) of **4b** as a light tan colored solid. ¹H NMR (CDCl₃, 400 MHz) (9 peaks) δ 1.1 (t, 3H, *J* = 7.0 Hz), 2.3 (s, 3H), 3.8 (s, 3H), 4.1 (q, 2H, *J* = 7.0 Hz), 5.3 (d, 1H), 5.8 (d, 1H), 6.8 (d, 2H, *J* = 8.2 Hz), 7.2 (d, 2H, *J* = 8.2 Hz), 8.3 (s, 1H); ¹³C NMR (CDCl₃, 100 MHz) (13 peaks) δ 14.1, 18.6, 55.1, 55.2, 60.0, 101.5, 113.9, 127.8, 136.0, 146.0, 153.4, 159.1, 165.7.

Reaction in BTF (entry 1b):

A mixture of *p*-anisaldehyde (0.501 g, 3.68 mmol), urea (0.243 g, 4.05 mmol, 1.1 equiv.) and ethyl acetoacetate (0.527 g, 0.512 mL, 4.05 mmol, 1.1 equiv.) in BTF (5.0 mL) was stirred at room temperature as iron(II) triflate (0.130 g, 0.368 mmol, 10.0 mol%) was added. The reaction mixture was heated at 70 °C (temperature-controlled hot plate). After 24 h, the mixture was cooled to room temperature. The

solvent was then removed by suction filtration and the resulting crude solid was triturated with CH₃OH/H₂O (1:1, v/v, 10.0 mL) twice to yield 0.632 g (59%) of **4b** as a light tan colored solid.

Reaction in 2-MeTHF (entry 1b):

A solution of *p*-anisaldehyde (0.504 g, 3.70 mmol, 1.0 eq), urea (0.244 g, 4.07 mmol, 1.1 eq) and ethyl acetoacetate (0.530 g, 0.519 mL, 4.07 mmol, 1.1 eq) in 2-MeTHF (6.0 mL) was stirred at room temperature as iron(II) triflate (0.131 g, 0.37 mmol, 10.0 mol%) was added. The reaction mixture was heated at 70 °C (temperature-controlled hot plate). After 24 h, the mixture was cooled to room temperature. The solvent was removed on a rotary evaporator, and the residue was triturated with CH₃OH/H₂O (1:1, v/v, 10.0 mL) twice to yield 0.787 g (74%) of **4b** as a light tan colored solid.

4. Conclusion

In summary, a new method for the Biginelli reaction that works in several different environmentally benign solvents, viz. isopropanol, benzotrifluoride, and 2-methyltetrahydrofuran, has been developed using an inexpensive and commercially available catalyst, iron(II) triflate. The best results were obtained in isopropanol, while 2-methyltetrahydrofuran is a viable option as it is a bio-derived solvent.

Acknowledgements

This material is based upon work supported by the National Science Foundation under CHE—1229133, which funded the purchase of a 400 MHz NMR spectrometer. RSM wishes to acknowledge an artistic and scholarly grant from Illinois Wesleyan University.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Atwal, K.S., Swanson, B.N., Unger, S.E., Floyd, D.M., Moreland, S., Hedberg, A., *et al.* (1991) Dihydropyrimidine Calcium Channel Blockers. 3. 3-Carbamoyl-4-aryl-1,2,3,4-tetrahydro-6-methyl-5-pyrimidinecarboxylic Acid Esters as Orally Effective Antihypertensive Agents. *Journal of Medicinal Chemistry*, **34**, 806-811. <https://doi.org/10.1021/jm00106a048>
- [2] Mayer, T.U., Kapoor, T.M., Haggarty, S.J., King, R.W., Schreiber, S.L. and Mitchison, T.J. (1999) Small Molecule Inhibitor of Mitotic Spindle Bipolarity Identified in a Phenotype-Based Screen. *Science*, **286**, 971-974. <https://doi.org/10.1126/science.286.5441.971>
- [3] Stefani, H.A., Oliveira, C.B., Almeida, R.B., Pereira, C.M.P., Braga, R.C., Cella, R., *et al.* (2006) Dihydropyrimidin-(2*H*)-ones Obtained by Ultrasound Irradiation: A New Class of Potential Antioxidant Agents. *European Journal of Medicinal Chemistry*, **41**, 513-518. <https://doi.org/10.1016/j.ejmech.2006.01.007>

- [4] Brands, M., Endermann, R., Gahlmann, R., Krüger, J. and Raddatz, S. (2003) Dihydropyrimidinones—A New Class of Anti-Staphylococcal Antibiotics. *Bioorganic & Medicinal Chemistry Letters*, **13**, 241-245. [https://doi.org/10.1016/s0960-894x\(02\)00880-6](https://doi.org/10.1016/s0960-894x(02)00880-6)
- [5] Biginelli, P. (1893) Derivati aldeidureidici degli eteri acetile dossal-acetico. *Gazzetta Chimica Italiana*, **23**, 360-413.
- [6] Touré, B.B. and Hall, D.G. (2009) Natural Product Synthesis Using Multicomponent Reaction Strategies. *Chemical Reviews*, **109**, 4439-4486. <https://doi.org/10.1021/cr800296p>
- [7] Tietze, L.F. (1996) Domino Reactions in Organic Synthesis. *Chemical Reviews*, **96**, 115-136. <https://doi.org/10.1021/cr950027e>
- [8] Kappe, C.O. (2000) Recent Advances in the Biginelli Dihydropyrimidine Synthesis. New Tricks from an Old Dog. *Accounts of Chemical Research*, **33**, 879-888. <https://doi.org/10.1021/ar000048h>
- [9] Kappe, C.O. (1997) A Reexamination of the Mechanism of the Biginelli Dihydropyrimidine Synthesis. Support for an *n*-Acyliminium Ion Intermediate. *The Journal of Organic Chemistry*, **62**, 7201-7204. <https://doi.org/10.1021/jo971010u>
- [10] Ahmed, B., Khan, R.A., Habibullah, and Keshari, M. (2009) An Improved Synthesis of Biginelli-Type Compounds via Phase-Transfer Catalysis. *Tetrahedron Letters*, **50**, 2889-2892. <https://doi.org/10.1016/j.tetlet.2009.03.177>
- [11] Chitra, S. and Pandiarajan, K. (2009) Calcium Fluoride: An Efficient and Reusable Catalyst for the Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones and Their Corresponding 2(1*H*)thione: An Improved High Yielding Protocol for the Biginelli Reaction. *Tetrahedron Letters*, **50**, 2222-2224. <https://doi.org/10.1016/j.tetlet.2009.02.162>
- [12] Nandurkar, N.S., Bhanushali, M.J., Bhor, M.D. and Bhanage, B.M. (2007) Y(NO₃)₃·6H₂O: A Novel and Reusable Catalyst for One Pot Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones under Solvent-Free Conditions. *Journal of Molecular Catalysis A: Chemical*, **271**, 14-17. <https://doi.org/10.1016/j.molcata.2007.02.021>
- [13] Adapa, S.R., Varala, R. and Alam, M.M. (2002) Bismuth Triflate Catalyzed One-Pot Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones: An Improved Protocol for the Biginelli Reaction. *Synlett*, No. 1, 0067-0070. <https://doi.org/10.1055/s-2003-36216>
- [14] Paraskar, A.S., Dewkar, G.K. and Sudalai, A. (2003) Cu(OTf)₂: A Reusable Catalyst for High-Yield Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones. *Tetrahedron Letters*, **44**, 3305-3308. [https://doi.org/10.1016/s0040-4039\(03\)00619-1](https://doi.org/10.1016/s0040-4039(03)00619-1)
- [15] Ranu, B.C., Hajra, A. and Jana, U. (2000) Indium(III) Chloride-Catalyzed One-Pot Synthesis of Dihydropyrimidinones by a Three-Component Coupling of 1,3-Dicarbonyl Compounds, Aldehydes, and Urea: An Improved Procedure for the Biginelli Reaction. *The Journal of Organic Chemistry*, **65**, 6270-6272. <https://doi.org/10.1021/jo000711f>
- [16] Bigi, F., Carloni, S., Frullanti, B., Maggi, R. and Sartori, G. (1999) A Revision of the Biginelli Reaction under Solid Acid Catalysis. Solvent-Free Synthesis of Dihydropyrimidines over Montmorillonite KSF. *Tetrahedron Letters*, **40**, 3465-3468. [https://doi.org/10.1016/s0040-4039\(99\)00424-4](https://doi.org/10.1016/s0040-4039(99)00424-4)
- [17] Tamaddon, F., Razmi, Z. and Jafari, A.A. (2010) Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones and 1,4-Dihydropyridines Using Ammonium Carbonate in Water. *Tetrahedron Letters*, **51**, 1187-1189. <https://doi.org/10.1016/j.tetlet.2009.12.098>
- [18] Debache, A., Amimour, M., Belfaitah, A., Rhouati, S. and Carboni, B. (2008) A One-Pot Biginelli Synthesis of 3,4-Dihydropyrimidin-2-(1*H*)-ones/thiones Catalyzed by

- Triphenylphosphine as Lewis Base. *Tetrahedron Letters*, **49**, 6119-6121. <https://doi.org/10.1016/j.tetlet.2008.08.016>
- [19] Shen, Z.-L., Xu, X.-P. and Ji, S.-J. (2010) Brønsted Base-Catalyzed One-Pot Three-Component Biginelli-Type Reaction: An Efficient Synthesis of 4,5,6-Triaryl-3,4-dihydropyrimidin-2(1*H*)-one and Mechanistic Study. *The Journal of Organic Chemistry*, **75**, 1162-1167. <https://doi.org/10.1021/jo902394y>
- [20] Pandey, J., Anand, N. and Tripathi, R.P. (2009) L-Proline Catalyzed Multicomponent Reaction of 3,4-Dihydro-(2*H*)-pyran, Urea/Thiourea, and Aldehydes: Diastereoselective Synthesis of Hexahydropyrano Pyrimidinones (Thiones). *Tetrahedron*, **65**, 9350-9356. <https://doi.org/10.1016/j.tet.2009.09.002>
- [21] Heravi, M.M., Moradi, R., Mohammadkhani, L. and Moradi, B. (2018) Current Progress in Asymmetric Biginelli Reaction: An Update. *Molecular Diversity*, **22**, 751-767. <https://doi.org/10.1007/s11030-018-9841-4>
- [22] Baruah, M.J., Dutta, R., Zaki, M.E.A. and Bania, K.K. (2024) Heterogeneous Iron-Based Catalysts for Organic Transformation Reactions: A Brief Overview. *Molecules*, **29**, Article No. 3177. <https://doi.org/10.3390/molecules29133177>
- [23] Lu, J. and Ma, H. (2000) Iron(III)-Catalyzed Synthesis of Dihydropyrimidinones: Improved Conditions for the Biginelli Reaction. *Synlett*, **2000**, 63-64. <https://doi.org/10.1055/s-2000-6469>
- [24] Krishna, T., Laxminarayana, E. and Kalita, D. (2020) FeF₃ as a Green Catalyst for the Synthesis of Dihydropyrimidines via Biginelli Reaction. *European Journal of Chemistry*, **11**, 206-212. <https://doi.org/10.5155/eurjchem.11.3.206-212.1992>
- [25] Adibi, H., Samimi, H.A. and Beygzadeh, M. (2007) Iron(III) Trifluoroacetate and Trifluoromethanesulfonate: Recyclable Lewis Acid Catalysts for One-Pot Synthesis of 3,4-Dihydropyrimidinones or Their Sulfur Analogues and 1,4-Dihydropyridines via Solvent-Free Biginelli and Hantzsch Condensation Protocols. *Catalysis Communications*, **8**, 2119-2124. <https://doi.org/10.1016/j.catcom.2007.04.022>
- [26] Starceovich, J.T., Laughlin, T.J. and Mohan, R.S. (2013) Iron(III) Tosylate Catalyzed Synthesis of 3,4-Dihydropyrimidin-2(1*H*)-ones/thiones via the Biginelli Reaction. *Tetrahedron Letters*, **54**, 983-985. <https://doi.org/10.1016/j.tetlet.2012.12.032>
- [27] Iron Fluoride (FeF₃). <https://pubchem.ncbi.nlm.nih.gov/compound/Iron-fluoride-FeF3#section=2D-Structure>
- [28] Phukan, M., Kalita, M.K. and Borah, R. (2010) A New Protocol for Biginelli (or like) Reaction under Solvent-Free Grinding Method Using Fe(NO₃)₃·9H₂O as Catalyst. *Green Chemistry Letters and Reviews*, **3**, 329-334. <https://doi.org/10.1080/17518253.2010.487841>
- [29] Rajni, Sahil, Raghav, N., Bendi, A. and Rao, G.B.D. (2026) Sustainable Biginelli Chemistry: Indian Contributions to Green Multicomponent Synthesis. *Discover Chemistry*, **3**, Article No. 172. <https://doi.org/10.1007/s44371-026-00582-z>
- [30] Mancheño, O.G., Dallimore, J., Plant, A. and Bolm, C. (2009) Iron(II) Triflate as an Efficient Catalyst for the Imination of Sulfoxides. *Organic Letters*, **11**, 2429-2432. <https://doi.org/10.1021/ol900660x>
- [31] Bonnamour, J. and Bolm, C. (2011) Iron(II) Triflate as a Catalyst for the Synthesis of Indoles by Intramolecular C-H Amination. *Organic Letters*, **13**, 2012-2014. <https://doi.org/10.1021/ol2004066>
- [32] Mayer, A.C., Salit, A. and Bolm, C. (2008) Iron-Catalysed Aziridination Reactions Promoted by an Ionic Liquid. *Chemical Communications*, No. 45, 5975-5977.

- <https://doi.org/10.1039/b813655f>
- [33] Kumawat, S. and Natte, K. (2024) Iron(II) Triflate as a Photocatalyst for Trifluoromethylation of Functionalized Arenes under Blue LED Light: Access to Bioactive Compounds. *Journal of Catalysis*, **434**, Article ID: 115506. <https://doi.org/10.1016/j.jcat.2024.115506>
- [34] Prat, D., Wells, A., Hayler, J., Sneddon, H., McElroy, C.R., Abou-Shehada, S., *et al.* (2016) CHEM21 Selection Guide of Classical- and Less Classical-Solvents. *Green Chemistry*, **18**, 288-296. <https://doi.org/10.1039/c5gc01008j>
- [35] Ogawa, A. and Curran, D.P. (1997) Benzotrifluoride: A Useful Alternative Solvent for Organic Reactions Currently Conducted in Dichloromethane and Related Solvents. *The Journal of Organic Chemistry*, **62**, 450-451. <https://doi.org/10.1021/jo9620324>
- [36] Maul, J.J., Ostrowski, P.J., Ublacker, G.A., Linclau, B. and Curran, D.P. (1999) Benzotrifluoride and Derivatives: Useful Solvents for Organic Synthesis and Fluorous Synthesis. In: Knochel, P., Ed., *Modern Solvents in Organic Synthesis*, Springer, 79-105. https://doi.org/10.1007/3-540-48664-x_4
- [37] Pace, V., Hoyos, P., Castoldi, L., Domínguez de María, P. and Alcántara, A.R. (2012) 2-Methyltetrahydrofuran (2-MeTHF): A Biomass-Derived Solvent with Broad Application in Organic Chemistry. *ChemSusChem*, **5**, 1369-1379. <https://doi.org/10.1002/cssc.201100780>
- [38] Fu, N.-Y., Yuan, Y.-F., Cao, Z., Wang, S.-W., Wang, J.-T. and Peppe, C. (2002) Indium(III) Bromide-Catalyzed Preparation of Dihydropyrimidinones: Improved Protocol Conditions for the Biginelli Reaction. *Tetrahedron*, **58**, 4801-4807. [https://doi.org/10.1016/s0040-4020\(02\)00455-6](https://doi.org/10.1016/s0040-4020(02)00455-6)
- [39] Li, P., Regati, S., Butcher, R.J., Arman, H.D., Chen, Z., Xiang, S., *et al.* (2011) Hydrogen-Bonding 2D Metal-Organic Solids as Highly Robust and Efficient Heterogeneous Green Catalysts for Biginelli Reaction. *Tetrahedron Letters*, **52**, 6220-6222. <https://doi.org/10.1016/j.tetlet.2011.09.099>
- [40] Ma, Y., Qian, C., Wang, L. and Yang, M. (2000) Lanthanide Triflate Catalyzed Biginelli Reaction. One-Pot Synthesis of Dihydropyrimidinones under Solvent-Free Conditions. *The Journal of Organic Chemistry*, **65**, 3864-3868. <https://doi.org/10.1021/jo9919052>
- [41] Ghosh, R., Maiti, S. and Chakraborty, A. (2004) In(OTf)₃-Catalysed One-Pot Synthesis of 3,4-Dihydropyrimidin-2(1H)-ones. *Journal of Molecular Catalysis A: Chemical*, **217**, 47-50. <https://doi.org/10.1016/j.molcata.2004.02.025>