

Association between Prepregnancy Obesity and Adverse Birth Outcomes among Mississippi Women

Ebele Okoye^{1,2*}, Vincent L. Mendy¹, Mary Shaw^{2,3}, Jung Hye Sung¹, Azad Bhuiyan¹, Marinelle Payton^{1,4}

¹Department of Epidemiology & Biostatistics, College of Health Sciences, Jackson State University, Jackson, MS, USA

²Mississippi Delta Center of Excellence in Maternal Health, Jackson, MS, USA

³Department of Behavioral & Environmental Health, College of Health Sciences, Jackson State University, Jackson, MS, USA

⁴Jackson Heart Study Graduate Training and Education Center, Jackson, MS, USA

Email: *ebele.c.okoye@students.jsums.edu, vincent.l.mendy@jsums.edu, mary.shaw@jsums.edu, jung.h.lee@jsums.edu,

azad.r.bhuiyan@jsums.edu, marinelle.payton@jsums.edu

How to cite this paper: Okoye, E., Mendy, V.L., Shaw, M., Sung, J.H., Bhuiyan, A. and Payton, M. (2026) Association between Prepregnancy Obesity and Adverse Birth Outcomes among Mississippi Women. *Open Journal of Obstetrics and Gynecology*, 16, 1572-1591.

<https://doi.org/10.4236/ojog.2026.1610141>

Received: September 17, 2026

Accepted: October 27, 2026

Published: October 30, 2026

Copyright © 2026 by author(s) and Scientific Research Publishing Inc.

This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Background: Prepregnancy obesity is associated with increased risk of adverse maternal and neonatal outcomes, including preterm birth and cesarean delivery. However, limited evidence exists from Mississippi-specific populations despite the state's high burden of obesity and adverse birth outcomes. **Objective:** This study examined the association between prepregnancy obesity and adverse birth outcomes, including cesarean delivery and preterm birth, among Mississippi women. **Methods:** We conducted a cross-sectional study using data from the Mississippi Pregnancy Risk Assessment Monitoring System, 2016-2022. The exposure was prepregnancy obesity (BMI ≥ 30.0 kg/m²), and the outcomes were cesarean delivery and preterm birth. Multivariable logistic regression analyses estimated the crude ratios (ORs) and adjusted odds ratios (AORs) with 95% confidence intervals (CIs), adjusting for covariates. **Results:** This study showed that among Mississippi women with live births (7,016), 51.6% were primarily White, and 53.4% were unmarried. About 57.3% had higher school education, 42.0% earned $\leq \$20,000$ annually, and 77.5% had insurance before pregnancy. There was a statistically significant association between prepregnancy obesity and cesarean delivery in all the models. Mississippi women with prepregnancy obesity had higher odds of cesarean delivery (AOR = 2.09; 95% CI: 1.86 - 2.42) compared to women without obesity, after adjusting for covariates. Although prepregnancy obesity was associated with preterm birth in the unadjusted analyses (OR = 1.29; 95% CI: 1.11 - 1.51), but the association was not statistically significant after adjusting for covariates (AOR = 1.10; 95%

CI: 0.93 - 1.31). **Conclusion:** Prepregnancy obesity is an independent predictor of cesarean delivery among women in Mississippi. These findings may inform public health practitioners and policymakers in developing interventions that promote preconception weight screening and obesity prevention.

Keywords

Adverse Pregnancy Outcomes, Cesarean Delivery, Maternal Health, Prepregnancy Obesity, Preterm Birth, PRAMS, Mississippi

1. Introduction

Obesity has emerged as one of the most significant public health challenges in the United States and disproportionately affects women of reproductive age [1]. Prepregnancy obesity, commonly defined as a body mass index (BMI) ≥ 30 kg/m² before conception, has increased substantially over the past several decades and is recognized as an important modifiable risk factor for adverse maternal and infant health outcomes [1] [2]. According to the Centers for Disease Control and Prevention (CDC), approximately 41.9% of U.S. adults had obesity during 2017 - 2020, with the highest prevalence observed among non-Hispanic Black adults and populations residing in the southern United States [3]. The rising prevalence of obesity among women entering pregnancy poses considerable challenges to maternal and child health and contributes substantially to healthcare expenditures and long-term morbidity [3].

Prepregnancy obesity has been associated with a wide range of adverse maternal and neonatal outcomes. Previous studies reported that women with prepregnancy obesity are at increased risk of gestational diabetes mellitus, gestational hypertension, preeclampsia, labor complications, cesarean delivery, postpartum hemorrhage, and maternal mortality [4]-[6]. Maternal obesity also adversely influences fetal growth and placental function and has been linked to several unfavorable birth outcomes, including preterm birth, macrosomia, congenital anomalies, stillbirth, neonatal intensive care unit admission, and infant mortality [4] [5] [7]. These adverse outcomes have significant implications not only for maternal and infant health during the perinatal period but also for long-term health trajectories, increasing the risk of obesity, metabolic disorders, and cardiovascular diseases among offspring later in life [8].

Among adverse birth outcomes, preterm birth and cesarean delivery remain major public health concerns in the United States, particularly in Mississippi. Preterm birth, defined as delivery before 37 completed weeks of gestation, is the leading cause of neonatal morbidity and mortality globally and is associated with respiratory distress syndrome, neurodevelopmental impairment, visual and hearing deficits, and chronic health conditions extending into adulthood [9] [10]. Similarly, although cesarean delivery is often a life-saving intervention when medically indicated, cesarean birth is associated with increased risks of maternal infection, hemorrhage, thromboembolic complications, prolonged hospitalization, and ad-

verse outcomes in subsequent pregnancies [11]. Previous studies have shown that women with obesity are more likely to undergo cesarean delivery due to obesity-related complications such as fetal macrosomia, dysfunctional labor, and pregnancy-related hypertensive disorders [5] [12]. Mississippi bears a disproportionate burden of obesity and poor maternal and infant health outcomes relative to many other states in the United States. In 2024, approximately 42.2% of women of childbearing age in Mississippi were classified as obese, exceeding national estimates [13]. Mississippi also consistently reports among the highest rates of preterm birth in the nation. According to the March of Dimes, 15.0% of live births in Mississippi in 2024 were preterm, and 37.9% were delivered by cesarean section [13] [14]. Furthermore, substantial racial and socioeconomic disparities persist, with Black women experiencing disproportionately higher rates of obesity, preterm birth, and adverse maternal outcomes compared with White women [14] [15]. These disparities are influenced by multiple social determinants of health, including poverty, limited access to healthcare, structural inequities, and chronic stress, indicating the need for targeted public health interventions [15].

Although associations between maternal obesity and adverse pregnancy outcomes are well established in the literature, much of the existing evidence is derived from nationally representative or non-state-specific populations, which may limit its applicability in Mississippi [5] [6] [12]. Population-based studies specifically examining the relationship between prepregnancy obesity and adverse birth outcomes among women in Mississippi remain limited. Given the state's persistently high prevalence of obesity and adverse birth outcomes, a clearer understanding of the impact of prepregnancy obesity on maternal and infant health is essential to inform targeted interventions and evidence-based maternal health policies.

Therefore, this study examines the association between prepregnancy obesity and adverse birth outcomes among Mississippi women with recent live births using data from the Pregnancy Risk Assessment Monitoring System (PRAMS), 2016-2022. So, the study assesses whether prepregnancy obesity is associated with higher odds of preterm birth and cesarean delivery after adjusting for key maternal sociodemographic characteristics. Findings from this study may inform preconception counseling and targeted maternal health interventions to reduce obesity-related risks and improve birth outcomes among women in Mississippi.

2. Materials and Methods

2.1. Study Design & Study Population

This study was part of a larger research project. We conducted a cross-sectional secondary analysis of quantitative data from the Mississippi PRAMS Phase 8 dataset (2016-2022). The Pregnancy Risk Assessment Monitoring System (PRAMS) is a surveillance project of the Centers for Disease Control and Prevention (CDC) in collaboration with state health departments. PRAMS collects state-specific, population-based data on maternal attitudes, behaviors, and experiences before, during, and shortly after pregnancy among women who recently had a live birth.

The study population comprises Mississippi resident women of reproductive age (18 - 44 years) who gave birth between 2016 and 2022 and completed the PRAMS survey. This study included women with and without prepregnancy obesity, with obesity defined as a body mass index (BMI) of $\geq 30 \text{ kg/m}^2$, in accordance with CDC guidelines.

Participants were eligible for inclusion if they were Mississippi residents who delivered a live singleton infant, had a recorded prepregnancy BMI, and completed the PRAMS Phase 8 questionnaire. Participants were subsequently classified as having prepregnancy obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$) or no prepregnancy obesity ($\text{BMI} < 30 \text{ kg/m}^2$). Women with multiple gestations (e.g., twins, triplets, or higher-order births) or reported preexisting diabetes or hypertension were excluded.

For each birth outcome, analyses were conducted among participants with available information on that outcome. Participants with missing values for variables included in a given regression model were excluded from that analysis. Consequently, the analytic sample size varied across outcomes and regression models based on the availability of the required variables.

2.2. Measures

The independent variable in this study was maternal obesity, defined as prepregnancy obesity. Prepregnancy body mass index (BMI) was calculated using self-reported PRAMS variables MAT_HTFT (height in feet), MAT_HTIN (height in inches), and MAT_PRWT (pregnancy weight), as described in the PRAMS codebook. Height was converted to meters and weight to kilograms, and BMI was calculated as weight in kilograms divided by height in meters squared (kg/m^2). Prepregnancy obesity was defined as a $\text{BMI} \geq 30 \text{ kg/m}^2$, whereas participants with a $\text{BMI} < 30 \text{ kg/m}^2$ were classified as non-obese.

Cesarean delivery was identified using the PRAMS variable DEL6HOW (Delivery-method) and categorized as a binary outcome (cesarean delivery vs. vaginal delivery). Gestational age was obtained from the PRAMS variable GEST_WK_NAPHSIS (Clinical estimate of gestational age). Preterm birth was defined as delivery before 37 completed weeks of gestation and categorized as a binary outcome (preterm vs. term). BMI was categorized according to the Centers for Disease Control and Prevention classification criteria. Women with a $\text{BMI} \geq 30 \text{ kg/m}^2$ were classified as obese, while women with a $\text{BMI} < 30 \text{ kg/m}^2$ were classified as non-obese. Prepregnancy obesity was operationalized as a binary variable (obese vs. non-obese [reference group]). The primary outcomes were adverse birth outcomes, including preterm birth and cesarean delivery. Each outcome was operationalized as a binary variable (Yes/No), indicating the presence or absence of the complication. The covariates were selected based on prior literature and their relevance to adverse birth outcomes. They included maternal age, race, marital status, educational attainment, household income, smoking before pregnancy, exercise before pregnancy, and insurance status.

2.3. Statistical Analysis

Descriptive statistics were used to summarize the characteristics of the study population, the prevalence of prepregnancy obesity, and the distribution of adverse birth outcomes. Frequencies and percentages were calculated as appropriate. Bivariate analyses were conducted to examine the associations between covariates and the two birth outcomes, cesarean delivery and preterm birth. Multivariable logistic regression models were used to assess the association between prepregnancy obesity and each adverse birth outcome, adjusting for potential covariates. Adjusted odds ratios (AORs) and 95% confidence intervals (CIs) were calculated.

Data Analysis

All statistical analyses were conducted using SAS version 9.4. Prior to analysis, Mississippi PRAMS datasets from 2016 through 2022 were merged to create a pooled cross-sectional analytic dataset. To account for multi-year pooling of independent cross-sectional samples, the survey (sampling) weights were adjusted by dividing the original sampling weight by seven (the number of survey years combined) to produce an average annual weight. Pooling PRAMS data across multiple years enabled a combined analysis of data from 2016 to 2022, accounting for the complex survey design and applying appropriate sample weights. Then, the dataset was prepared to account for the complex survey design by incorporating the sampling weight (WTANAL) and stratification variable (STRATUMC). SAS survey procedures were used to appropriately adjust for weighting and stratification. Weighted descriptive statistics were generated using PROC SURVEYFREQ to summarize the demographic characteristics of the study population. Weighted cross-tabulations were used to describe the distribution and weighted prevalence of cesarean delivery and preterm birth across maternal characteristics. Rao-Scott chi-square tests were used to assess differences in the distribution of each adverse pregnancy outcome across categories of maternal characteristics. Logistic regression analyses were performed using PROC SURVEYLOGISTIC to estimate odds ratios and 95% confidence intervals (CIs) for the association between prepregnancy obesity and adverse birth outcomes, including preterm birth and cesarean delivery.

Unadjusted models were first estimated to assess crude associations between prepregnancy obesity and adverse birth outcomes. Subsequently, multivariable logistic regression models were fitted to control for potential confounders. Three models were specified: Model 1 was unadjusted; Model 2 was adjusted for maternal age and race; and Model 3 was further adjusted for maternal age, race, marital status, education, income level, insurance status, smoking, and exercise. Odds ratios (ORs) and adjusted odds ratios (AORs) with corresponding 95% confidence intervals (CIs) were reported to quantify the associations between prepregnancy obesity and adverse birth outcomes among Mississippi women of reproductive age. Statistical significance was determined using a two-sided alpha level of 0.05, and associations were considered statistically significant if the p-value was <0.05 and the 95% confidence interval did not include 1.0.

2.4. Ethical Considerations

This study used secondary, de-identified data from the Mississippi Pregnancy Risk Assessment Monitoring System (PRAMS). No direct contact with participants or identifiable personal information was involved. The study was reviewed and approved by the Jackson State University Institutional Review Board and did not require informed consent from individual participants.

3. Results

3.1. Sociodemographic and Adverse Birth Outcome Characteristics of the Study Population

This study comprised 7016 participants, which, when weighted using the complex design, totals 33,734 and represents the average annual population of Mississippi women, accounting for those who may have been undersampled or oversampled during the survey. **Table 1** summarizes the sociodemographic and adverse birth outcomes characteristics of Mississippi women with recent live births who participated in and completed the PRAMS survey from 2016 to 2022. Most (32.4%) women aged 25 - 29 years represent the largest age group, followed closely by those aged ≥ 30 years (31.1%). Women aged 20 - 24 years accounted for 28.1% of the population, while adolescents aged <20 years comprised a smaller proportion of the sample (8.4%). Regarding race, White women represented just over half (51.6%) of the study population, Black women accounted for 44.9%, while other races accounted for 6.4%. More than half (53.4%) of the women were not married at the time of pregnancy, while 46.6% were married. Educational attainment varied widely. Approximately 12.1% of women had less than a high school education, and 30.6% had a high school education. However, a notable proportion, over half (57.3%) of the women had a higher level of education than high school. A large proportion of women reported low household income, with 42.0% earning $\leq \$20,000$ annually. Approximately 24.4% earned between \$20,001 and \$40,000, while 21.5% earned \$60,000 or more. Regarding insurance, most women (77.5%) reportedly had insurance before pregnancy, while 22.5% did not. The majority (91.2%) of the women reported not smoking before pregnancy; however, 8.8% reported smoking prior to pregnancy. Approximately 34.5% of women reported exercising before pregnancy, whereas 65.5% did not.

Table 1. Characteristics of the study population, PRAMS 2016-2022.

Characteristics	n ^a	N ^b	% ^c	95% CI
Age (Years)				
<20	620	2824	8.4	7.5 - 9.3
20 - 24	1948	9489	28.1	26.7 - 29.6
25 - 29	2159	10,923	32.4	30.9 - 33.9
≥ 30	2289	10,498	31.1	29.7 - 32.6

Continued

Educational level				
<High school	905	4072	12.1	11.0 - 13.2
High school	2150	10,301	30.6	29.1 - 32.1
>High school	3942	19,285	57.3	55.7 - 58.9
Race				
White	3134	17,391	51.6	50.0 - 53.2
Black	3545	14,179	44.9	40.5 - 43.6
Other race	337	2164	6.4	5.6 - 7.3
Marital status				
Married	2890	15,711	46.6	45.0 - 48.2
Not married	4123	18,020	53.4	51.8 - 55.0
Annual household income, \$				
≤20,000	3082	13,164	42.0	40.4 - 43.7
>20,000 to ≤40,000	1558	7632	24.4	22.9 - 25.8
>40,000 to ≤60,000	687	3793	12.1	11.0 - 13.2
>60,000	1154	6727	21.5	20.1 - 22.9
Insurance status before pregnancy				
Uninsured	1553	7581	22.5	21.2 - 23.9
Insured	5453	26,128	77.5	76.2 - 78.9
Smoking before pregnancy				
No	6265	30,750	91.2	90.4 - 92.1
Yes	742	2955	8.8	7.9 - 9.6
Prepregnancy exercise				
Yes	2293	11,574	34.5	32.9 - 36.0
No	4685	21,997	65.5	63.9 - 67.1
Prepregnancy obesity				
Yes	2747	13,183	39.1	37.5 - 40.6
No	4269	20,551	60.9	59.4 - 62.5
Delivery method				
Vaginal	3716	20,441	62.7	61.1 - 64.2
Caesarean section	3081	12,184	37.3	35.8 - 38.9

Continued

Preterm birth				
Yes	2740	4372	13.0	12.2 - 13.8
No	4268	29,332	87.0	86.2 - 87.8

Note: n^a = Unweighted number; N^b = Weighted number; %^c = Weighted percentage; CI = Confidence Interval; PRAMS Pregnancy Risk Assessment Monitoring System.

Regarding adverse birth outcomes, 39.1% reported prepregnancy obesity, while 60.9% were not obese prior to pregnancy. In terms of delivery method, vaginal delivery was the most common birth method, occurring in 62.7% of women, while 37.3% had cesarean delivery. Preterm birth (<37 weeks) occurred in 13.0% of women during pregnancy, while 87.0% delivered at term.

3.2. Maternal Characteristics by Adverse Birth Outcome among Mississippi Women

3.2.1. Prevalence of Preterm Birth by Maternal Characteristics among Mississippi Women

Among Mississippi women with recent live births, the overall weighted prevalence of preterm birth was 13.0% (95% CI: 12.2% - 13.8%), **Table 2**. Preterm birth prevalence was significantly higher among women with prepregnancy obesity (14.7%) than among those without obesity (11.8%; $p = 0.001$). Significant differences in preterm birth prevalence were also observed by maternal education ($p = 0.036$), race ($p < 0.001$), marital status ($p < 0.001$), and household income ($p < 0.001$). Preterm birth prevalence was highest among women with a high school education (14.4%), Black women (16.7%), unmarried women (14.5%), and women with annual household income $\leq$ \$20,000 (15.4%).

Table 2. Prevalence of preterm birth by maternal characteristics among Mississippi women, PRAMS 2016-2022.

Preterm Birth				
Maternal characteristics	n ^a	N ^b	% ^c (95% CI)	p-value
Prepregnancy obesity				0.001
Not obese	1572	4238	11.8 (10.9 - 12.8)	
Obese	1149	2736	14.7 (13.3 - 16.3)	
Maternal age at delivery				0.154
<20 years	244	618	11.0 (8.9 - 13.4)	
20 - 24 years	771	1946	13.1 (11.6 - 14.8)	
25 - 29 years	801	2157	12.3 (10.9 - 13.9)	
≥ 30 years	924	2287	14.1 (12.7 - 15.7)	

Continued

Maternal education			0.036
<High school	364	902	14.0 (11.6 - 16.7)
High school	905	2149	14.4 (12.8 - 16.0)
>High school	1462	3941	12.0 (11.0 - 13.1)
Maternal race			<0.001
White	993	3130	10.3 (9.3 - 11.5)
Black	1653	3541	16.7 (15.4 - 18.1)
Other races	94	337	9.5 (6.7 - 13.3)
Marital status			<0.001
Not married	1767	4117	14.5 (13.4 - 15.7)
Married	972	2890	11.2 (10.1 - 12.5)
Annual household income\$			<0.001
≤20,000	1359	3077	15.4 (14.0 - 16.8)
>20,000 - ≤40,000	597	1556	12.8 (11.2 - 14.6)
>40,000 - \$60,000	234	686	11.6 (9.4 - 14.3)
≥\$60,000	346	1154	9.8 (8.1 - 11.8)
Insurance status before pregnancy			0.142
Uninsured	571	1549	11.8 (10.2 - 13.6)
Insured	2162	5449	13.3 (12.4 - 14.3)
Smoking before pregnancy			0.293
No	2422	6261	12.8 (12.0 - 13.7)
Yes	310	732	14.3 (11.8 - 17.1)
Prepregnancy exercise			0.500
No	1868	4679	13.1 (12.2 - 14.2)
Yes	855	2291	12.5 (11.1 - 14.1)

Note: n^a = Unweighted number of women who reported gestational hypertension/preeclampsia; N^b = the total unweighted number of women in each maternal characteristic category; %^c = Percentage; CI = Confidence interval of the survey-weighted; p-value = <0.05; PRAMS Pregnancy Risk Assessment Monitoring System.

In contrast, preterm birth prevalence did not differ significantly by maternal age ($p = 0.154$), insurance status ($p = 0.142$), smoking before pregnancy ($p = 0.293$), or prepregnancy exercise ($p = 0.500$). Among women who reported prepregnancy exercise ≥ 3 days per week, the prevalence of preterm birth was 13.1%, compared with 12.5% among those who did not engage in prepregnancy exercise.

3.2.2. Prevalence of Cesarean Delivery by Maternal Characteristics among Mississippi Women

The weighted prevalence of cesarean delivery among Mississippi women with recent live births was 37.3%. The prevalence varied significantly by several maternal characteristics (**Table 3**). Women with prepregnancy obesity had a substantially higher prevalence of cesarean delivery than women who were not obese (48.5% vs. 30.1%; $p < 0.001$).

Table 3. Prevalence of cesarean delivery by maternal characteristics among Mississippi women, PRAMS 2016-2022.

Maternal characteristic	Cesarean Delivery			p-value
	n ^a	N ^b	% ^c (95% CI)	
Prepregnancy obesity				<0.001
Not obese	1588	4102	30.1 (28.3 - 32.0)	
Obese	1476	2656	48.5 (46.0 - 51.1)	
Maternal age at delivery				<0.001
<20 years	167	593	20.4 (16.2 - 25.2)	
20 - 24 years	766	1886	31.6 (28.9 - 34.5)	
25 - 29 years	992	2099	40.6 (37.8 - 43.5)	
≥30 years	1156	2219	43.6 (40.9 - 46.3)	
Maternal education				<0.001
<High school	328	861	29.7 (25.6 - 34.2)	
High school	913	2069	35.6 (32.8 - 38.5)	
>High school	1836	3850	39.8 (37.8 - 41.9)	
Maternal race				0.206
White	1351	3067	37.0 (34.8 - 39.2)	
Black	1606	3405	38.5 (36.2 - 40.9)	
Other races	124	325	32.6 (26.6 - 39.2)	
Marital status				0.011
Not married	1746	3969	35.4 (33.4 - 37.6)	
Married	1335	2825	39.5 (37.2 - 41.8)	
Annual household income, \$				0.361
≤20,000	1356	3042	35.7 (33.3 - 38.2)	
>20,000 - ≤40,000	730	1543	38.2 (35.0 - 41.5)	
>40,000 - ≤60,000	319	679	39.5 (34.8 - 44.4)	
≥60,000	514	1134	38.6 (35.2 - 42.1)	

Continued

Insurance status before pregnancy			0.707
Uninsured	669	1506	36.8 (33.6 - 40.1)
Insured	2409	5281	37.5 (35.8 - 39.3)
Smoking before pregnancy			0.498
No	2744	6066	37.1 (35.5 - 38.8)
Yes	329	714	39.0 (34.0 - 44.1)
Prepregnancy exercise			0.566
Yes	1018	2217	37.9 (35.3 - 40.6)
No	2044	4543	37.0 (35.1 - 38.9)

Note: n^a = Unweighted number of women who reported gestational hypertension/preeclampsia; N^b = the total unweighted number of women in each maternal characteristic category; %^c = Percentage; CI = Confidence interval of the survey-weighted; p-value = <0.05; PRAMS Pregnancy Risk Assessment Monitoring System.

Cesarean delivery prevalence also differed significantly by maternal age, marital status, and education. The prevalence increased with age, from 20.4% among women younger than 20 years to 43.6% among women aged ≥ 30 years ($p < 0.001$). Married women had a higher prevalence of cesarean delivery than unmarried women (39.5% vs. 35.4%; $p = 0.011$). Similarly, cesarean delivery prevalence increased with educational attainment, from 29.7% among women with less than a high school education to 39.8% among those with more than a high school education ($p < 0.001$).

No statistically significant differences in the prevalence of cesarean delivery were observed by maternal race ($p = 0.206$), annual household income ($p = 0.361$), insurance status before pregnancy ($p = 0.707$), smoking before pregnancy ($p = 0.498$), or prepregnancy exercise ($p = 0.566$). The prevalence of cesarean delivery was 37.9% among women who exercised before pregnancy compared with 37.0% among women who did not exercise.

3.3. Odds of Adverse Birth Outcomes and Prepregnancy Obesity among Mississippi Women, PRAMS 2016-2022

Table 4 presents the association between prepregnancy obesity and cesarean delivery among Mississippi women who participated in PRAMS from 2016 to 2022. In the unadjusted model (Model 1), women with prepregnancy obesity had more than twice the odds of delivering by cesarean section compared with women who were not obese prior to pregnancy (OR = 2.18; 95% CI: 1.91 - 2.50). After adjustment for maternal age and race (Model 2), the association remained statistically significant, with obese women having 2.08 times higher odds of cesarean delivery (AOR = 2.08; 95% CI: 1.80 - 2.40). In the fully adjusted model (Model 3), which controlled for covariates including sociodemographic factors (maternal age, race, marital status, education, income, and insurance) and behavioral factors (smoking and exercise), prepregnancy obesity remained independently associated with

cesarean delivery (AOR = 2.09; 95% CI: 1.86 - 2.42).

Table 4. Odds of cesarean delivery and prepregnancy obesity among Mississippi women, PRAMS 2016-2022.

Characteristic		Model 1	Model 2	Model 3
Outcome	Prepregnancy obesity	OR ^a (95% CI)	AOR ^b (95% CI)	AOR ^c (95% CI)
Cesarean delivery	Non-obese	1.00 (Reference)	1.00 (Reference)	1.00 (Reference)
	Obese	2.18 (1.91 - 2.50)	2.08 (1.80 - 2.40)	2.09 (1.86 - 2.42)
	Unweighted (n)	6758	6758	6331
	Weighted (N)	32,467	32,467	30,571

Note: AOR: adjusted odds ratio; PRAMS, Pregnancy Risk Assessment Monitoring System; CI: confidence interval; OR: odds ratio; ^aunadjusted; ^badjusted for age, race; ^cadjusted for age, race, marital status, education, income, insurance, smoking, exercise. Non-obese (BMI <30 kg/m²) was the reference category; obese was defined as BMI ≥30 kg/m². Analytic sample sizes vary across models because participants with missing values for variables included in a given model were excluded from that analysis.

Table 5 presents the odds of preterm birth (<37 weeks of gestation) in relation to prepregnancy obesity among Mississippi women participating in PRAMS from 2016 to 2022. In the unadjusted model (Model 1), women with prepregnancy obesity had 29% higher odds of preterm birth compared with women without obesity (OR = 1.29; 95% CI: 1.11 - 1.51). After adjustment for maternal age and race (Model 2), the association was attenuated and no longer statistically significant (AOR = 1.14; 95% CI: 0.97 - 1.34). After further adjustment for covariates, including maternal age, race, marital status, education, household income, insurance status, smoking, and exercise before pregnancy (Model 3), the association remained non-significant (AOR = 1.10; 95% CI: 0.93 - 1.31).

Table 5. Odds of preterm birth and prepregnancy obesity among Mississippi women, PRAMS 2016-2022.

Characteristic		Model 1	Model 2	Model 3
Outcome	Prepregnancy obesity	OR ^a (95% CI)	AOR ^b (95% CI)	AOR ^c (95% CI)
Preterm birth (<37 weeks of gestation)	Non-obese	1.00 (Reference)	1.00 (Reference)	1.00 (Reference)
	Obese	1.29 (1.11 - 1.51)	1.14 (0.97 - 1.34)	1.10 (0.93 - 1.31)
	Unweighted (n)	6974	6974	6411
	Weighted (N)	33,549	33,549	31,020

Note: AOR: adjusted odds ratio; PRAMS, Pregnancy Risk Assessment Monitoring System; CI: confidence interval; OR: odds ratio; ^aunadjusted; ^badjusted for age, race; ^cadjusted for age, race, marital status, education, income, insurance, smoking, exercise. Non-obese (BMI <30 kg/m²) was the reference category; obese was defined as BMI ≥30 kg/m². Analytic sample sizes vary across models because participants with missing values for variables included in a given model were excluded from that analysis.

4. Discussion

The study population experienced a substantial burden of socioeconomic and maternal health risk factors. Prepregnancy obesity affected approximately 39% of

women, while cesarean delivery and preterm birth occurred in more than one-third and approximately 13% of births, respectively. Nearly half of the women reported annual household incomes below \$20,000, and a substantial proportion reported limited prepregnancy exercise, showing the importance of addressing socioeconomic conditions and modifiable maternal health behaviors in preconception care [16]-[25].

The weighted prevalence of cesarean delivery differed significantly by maternal age, marital status, educational attainment, and prepregnancy obesity. The higher prevalence among women aged ≥ 30 years is consistent with evidence linking advanced maternal age to increased cesarean delivery, potentially due to greater risk of pregnancy complications and labor-related indications [26]-[28]. Differences by marital status and education may reflect underlying differences in maternal characteristics, healthcare utilization, and social circumstances [29]-[31]. The substantially higher prevalence of cesarean delivery among women with prepregnancy obesity is consistent with previous evidence linking maternal obesity to cesarean birth [5] [6] [12] [32].

The weighted prevalence of preterm birth differed significantly by maternal race, marital status, educational attainment, household income, and prepregnancy obesity. The higher prevalence among Black women is consistent with persistent racial disparities in preterm birth in the United States, which are largely attributed to social, structural, and environmental factors rather than race itself [14] [16] [33]. Differences by marital status, education, and household income may similarly reflect disparities in social support, economic resources, healthcare access, and psychosocial stress [16] [34]. Women with prepregnancy obesity also had a higher prevalence of preterm birth, consistent with previous evidence linking maternal obesity with preterm delivery [5] [6] [12].

4.1. Association between Prepregnancy Obesity and Cesarean Delivery among Mississippi Women

This study found a strong and consistent association between prepregnancy obesity and cesarean delivery among Mississippi women with recent live births. Women with prepregnancy obesity had more than twice the odds of cesarean delivery compared with women without obesity, and this association remained statistically significant after adjustment for maternal age, race, sociodemographic characteristics, and behavioral factors. The persistence of the association across the regression models indicates that prepregnancy obesity was independently associated with cesarean delivery in this population. These findings are consistent with previous studies showing that maternal obesity is associated with an increased likelihood of cesarean delivery [5] [6] [35] [36].

Marchi *et al.* [5] and Vats *et al.* [6] reported significantly greater risks of cesarean delivery among women with prepregnancy obesity, while Rubens *et al.* [35], using a large U.S. population-based dataset, found approximately 70% higher odds of cesarean delivery among women with obesity. Similarly, Zhang *et al.* [36]

reported increased odds of cesarean delivery among women with overweight or obesity. The consistency of these findings across different populations and study designs supports the association observed in the present study.

Although the mechanisms underlying this association were not examined in the present study, several biological and obstetric pathways may contribute to the increased likelihood of cesarean delivery among women with prepregnancy obesity. Maternal obesity has been associated with pregnancy complications, including gestational diabetes and hypertensive disorders, as well as labor-related factors such as prolonged labor, less successful cervical ripening, and increased need for labor induction and augmentation, which may increase the likelihood of operative delivery [36] [37]. Therefore, the observed association may reflect, in part, the greater occurrence of obesity-related pregnancy and intrapartum complications among women with prepregnancy obesity.

These findings show the importance of addressing obesity before conception as part of comprehensive maternal healthcare. Preconception counseling, weight management, and evidence-based lifestyle interventions may help optimize maternal health and reduce obesity-related pregnancy complications, including the risk of cesarean delivery [38]–[40].

4.2. Association between Prepregnancy Obesity and Preterm Birth among Mississippi Women

We found that prepregnancy obesity was associated with increased odds of preterm birth in the unadjusted analysis; however, the association was no longer statistically significant after adjustment for maternal age, race, and other sociodemographic and behavioral characteristics. These findings suggest that maternal demographic, socioeconomic, and behavioral characteristics may partly account for the crude association observed between prepregnancy obesity and preterm birth.

The attenuation of the association after adjustment is consistent with the multifactorial etiology of preterm birth, reflecting a complex interplay among biological, behavioral, and social determinants of health [41]. Maternal socioeconomic disadvantage, smoking, healthcare access, and environmental conditions have been associated with preterm birth and may contribute to the risk inequalities [41] [42]. Racial disparities in preterm birth may likewise reflect broader social and structural conditions rather than race itself [33]. These factors may be particularly relevant in Mississippi, where racial and socioeconomic disparities in maternal health remain substantial.

Our findings differ from several previous studies reporting an independent association between prepregnancy obesity and preterm birth. Marchi *et al.* [5] and Vats *et al.* [6] found that maternal obesity was associated with an elevated risk of preterm birth. More recently, Cornish *et al.* [43], using data from nearly 29 million pregnancies, found that higher maternal prepregnancy BMI was associated with increased risk of overall and medically indicated preterm birth after adjustment, although the associations varied by preterm birth phenotype and parity. Differ-

ences in study populations, outcome definitions, covariate adjustment, and regional characteristics may partly explain the discrepancy between these findings and those of the present study.

Although prepregnancy obesity was not independently associated with preterm birth after adjustment, the findings show the importance of considering maternal sociodemographic and behavioral characteristics when examining the risk of preterm birth. Strategies to reduce preterm birth in Mississippi should therefore extend beyond obesity prevention and incorporate approaches that address social determinants of health, improve access to quality preconception and prenatal care, reduce tobacco use, and promote maternal health equity [33] [41] [42].

5. Conclusion

This study examined the association between prepregnancy obesity and adverse birth outcomes among Mississippi women using PRAMS data from 2016 to 2022. Among Mississippi women, prepregnancy obesity was highly prevalent and was significantly associated with increased odds of cesarean delivery after adjusting for covariates, indicating an independent association between prepregnancy obesity and cesarean delivery. However, there was no association between prepregnancy obesity and preterm birth among Mississippi women with recent live births. Marked sociodemographic disparities were observed, with a higher prevalence of prepregnancy obesity among older, Black, unmarried women and those with lower household income, indicating important social and structural disparities in maternal health. These findings indicate the need for targeted, equity-focused preconception and prenatal interventions.

6. Study Implications

This study provides important implications for maternal and public health practice in Mississippi. The observed association between prepregnancy obesity and cesarean delivery indicates the need for strengthened care during preconception, which includes routine screening for obesity and early identification of women at elevated risk for pregnancy complications. Integrating weight management counseling and preventive services into primary care and reproductive health settings may help reduce obesity-related adverse birth outcomes.

The identified sociodemographic disparities suggest that interventions should be equity-focused and tailored to populations disproportionately affected by prepregnancy obesity, particularly among older women, Black women, unmarried women, and those with lower household income. Addressing these disparities requires not only individual-level behavioral interventions but also broader structural strategies to improve access to healthy foods, safe environments for physical activity, and affordable healthcare services. At the policy level, these findings support the development of state-level maternal health initiatives tailored to Mississippi women, prioritizing preconception health promotion and obesity prevention as key strategies to improve pregnancy outcomes. Therefore, this study shows

the importance of early, targeted, and multi-level interventions to reduce the burden of prepregnancy obesity and improve maternal and infant health outcomes in Mississippi.

7. Strengths and Limitations

A key strength of this study is the use of PRAMS data, a large, population-based surveillance system with robust weighting procedures that enhance the generalizability of findings to women with recent live births in Mississippi. Using multiple years of data (2016–2022) improves statistical power and yields more stable estimates. PRAMS data also provide important state-level surveillance, helping to characterize Mississippi's burden of maternal obesity and the associated risk of adverse birth outcomes. Additionally, the study accounted for the complex survey design and adjusted for relevant sociodemographic and health-related covariates, strengthening the validity of the observed associations. However, this study has some limitations. The cross-sectional design limits the ability to establish causality between prepregnancy obesity and adverse birth outcomes. Self-reported data on prepregnancy weight and height may introduce recall or reporting bias, potentially affecting BMI classification. In addition, information on parity and prior cesarean delivery was not available in the analytic dataset and, therefore, could not be included in the regression analysis. Because these factors may influence the likelihood of cesarean delivery, their potential confounding effects could not be assessed.

8. Future Directions

Further research should explore the mechanisms linking prepregnancy obesity to adverse birth outcomes, including biological, behavioral, and healthcare-related factors. Also, longitudinal studies are needed to better establish the temporal relationships and causal mechanisms between prepregnancy obesity and adverse birth outcomes. Finally, future work should evaluate the effectiveness of targeted preconception and prenatal interventions aimed at reducing obesity prevalence and improving maternal and infant health outcomes, particularly among disproportionately affected populations.

Acknowledgements

The authors gratefully acknowledge the Mississippi State Department of Health (MSDH) for making the Pregnancy Risk Assessment Monitoring System (PRAMS) data available for this study.

Author Contributions

Conceptualization, E.O.; methodology, E.O.; software, E.O.; validation, E.O., V.L.M., A.B., J.L., M.S., and M.P.; formal analysis, E.O., and V.L.M.; data curation, E.O.; writing—original draft preparation, E.O.; writing—review and editing, E.O., V.L.M., A.B., J.L., M.S., and M.P.; visualization, E.O.; supervision, V.L.M., A.B., J.L.,

M.S., and M.P. All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest regarding the publication of this paper.

References

- [1] Emmerich, S.D., Fryar, C.D., Stierman, B. and Ogden, C.L. (2024) Obesity and Severe Obesity Prevalence in Adults: United States, August 2021-August 2023. NCHS Data Brief (508).
- [2] Driscoll, A.K. and Gregory, E.C.W. (2020) Increases in Prepregnancy Obesity: United States, 2016-2019. NCHS Data Brief 392. <https://www.cdc.gov/nchs/products/index.htm>
- [3] Hales, C.M., Carroll, M.D., Fryar, C.D. and Ogden, C.L. (2024) Adult Obesity Facts. Centers for Disease Control and Prevention. <https://www.cdc.gov/obesity/adult-obesity-facts/index.html>
- [4] Aune, D., Saugstad, O.D., Henriksen, T. and Tonstad, S. (2014) Maternal Body Mass Index and the Risk of Fetal Death, Stillbirth, and Infant Death: A Systematic Review and Meta-Analysis. *JAMA*, **311**, 1536-1546. <https://doi.org/10.1001/jama.2014.2269>
- [5] Marchi, J., Berg, M., Dencker, A., Olander, E.K. and Begley, C. (2015) Risks Associated with Obesity in Pregnancy, for the Mother and Baby: A Systematic Review of Reviews. *Obesity Reviews*, **16**, 621-638. <https://doi.org/10.1111/obr.12288>
- [6] Vats, H., Saxena, R., Sachdeva, M.P., Walia, G.K. and Gupta, V. (2021) Impact of Maternal Pre-Pregnancy Body Mass Index on Maternal, Fetal and Neonatal Adverse Outcomes in the Worldwide Populations: A Systematic Review and Meta-Analysis. *Obesity Research & Clinical Practice*, **15**, 536-545. <https://doi.org/10.1016/j.orcp.2021.10.005>
- [7] Cnattingius, S., Villamor, E., Johansson, S., Bonamy, A.E., Persson, M., Wikström, A., et al. (2013) Maternal Obesity and Risk of Preterm Delivery. *JAMA*, **309**, 2362-2370. <https://doi.org/10.1001/jama.2013.6295>
- [8] Godfrey, K.M., Reynolds, R.M., Prescott, S.L., Nyirenda, M., Jaddoe, V.W.V., Eriksson, J.G., et al. (2017) Influence of Maternal Obesity on the Long-Term Health of Offspring. *The Lancet Diabetes & Endocrinology*, **5**, 53-64. [https://doi.org/10.1016/s2213-8587\(16\)30107-3](https://doi.org/10.1016/s2213-8587(16)30107-3)
- [9] Blencowe, H., Cousens, S., Chou, D., Oestergaard, M., Say, L., Moller, A., et al. (2013) Born Too Soon: The Global Epidemiology of 15 Million Preterm Births. *Reproductive Health*, **10**, Article No. S2. <https://doi.org/10.1186/1742-4755-10-s1-s2>
- [10] Walani, S.R. (2020) Global Burden of Preterm Birth. *International Journal of Gynecology & Obstetrics*, **150**, 31-33. <https://doi.org/10.1002/ijgo.13195>
- [11] Sandall, J., Tribe, R.M., Avery, L., Mola, G., Visser, G.H., Homer, C.S., et al. (2018) Short-Term and Long-Term Effects of Caesarean Section on the Health of Women and Children. *The Lancet*, **392**, 1349-1357. [https://doi.org/10.1016/s0140-6736\(18\)31930-5](https://doi.org/10.1016/s0140-6736(18)31930-5)
- [12] Krogh, L.Q., Glavind, J., Henriksen, T.B., Thornton, J., Fuglsang, J. and Boie, S. (2023) Full-Term Induction of Labor vs Expectant Management and Cesarean Delivery in Women with Obesity: Systematic Review and Meta-Analysis. *American Journal of Obstetrics & Gynecology MFM*, **5**, Article ID: 100909. <https://doi.org/10.1016/j.ajogmf.2023.100909>

- [13] March of Dimes (2026) Obesity among Women of Childbearing Age: United States, 2013-2024. PeriStats.
<https://www.marchofdimes.org/peri-stats/data?reg=99&top=17&stop=350&slev=1&obj=1>
- [14] March of Dimes (2024) PeriStats: Mississippi Data. March of Dimes.
<https://www.marchofdimes.org/peristats/state-summaries/mississippi>
- [15] Howell, E.A. (2018) Reducing Disparities in Severe Maternal Morbidity and Mortality. *Clinical Obstetrics & Gynecology*, **61**, 387-399.
<https://doi.org/10.1097/grf.0000000000000349>
- [16] Braveman, P.A., Heck, K., Egerter, S., Marchi, K.S., Dominguez, T.P., Cubbin, C., *et al.* (2015) The Role of Socioeconomic Factors in Black-White Disparities in Preterm Birth. *American Journal of Public Health*, **105**, 694-702.
<https://doi.org/10.2105/ajph.2014.302008>
- [17] Braveman, P., Egerter, S. and Williams, D.R. (2011) The Social Determinants of Health: Coming of Age. *Annual Review of Public Health*, **32**, 381-398.
<https://doi.org/10.1146/annurev-publhealth-031210-101218>
- [18] Wang, E., Glazer, K.B., Howell, E.A. and Janevic, T.M. (2020) Social Determinants of Pregnancy-Related Mortality and Morbidity in the United States. *Obstetrics & Gynecology*, **135**, 896-915. <https://doi.org/10.1097/aog.00000000000003762>
- [19] Declercq, E., MacDorman, M., Osterman, M., Belanoff, C. and Iverson, R. (2015) Pre-pregnancy Obesity and Primary Cesareans among Otherwise Low-Risk Mothers in 38 U.S. States in 2012. *Birth*, **42**, 309-318. <https://doi.org/10.1111/birt.12201>
- [20] Kominniarek, M.A. and Peaceman, A.M. (2017) Gestational Weight Gain. *American Journal of Obstetrics and Gynecology*, **217**, 642-651.
<https://doi.org/10.1016/j.ajog.2017.05.040>
- [21] McAuliffe, F.M., Killeen, S.L., Jacob, C.M., Hanson, M.A., Hadar, E., McIntyre, H.D., *et al.* (2020) Management of Prepregnancy, Pregnancy, and Postpartum Obesity from the FIGO Pregnancy and Non-Communicable Diseases Committee: A FIGO (International Federation of Gynecology and Obstetrics) Guideline. *International Journal of Gynecology & Obstetrics*, **151**, 16-36. <https://doi.org/10.1002/ijgo.13334>
- [22] Stephenson, J., Heslehurst, N., Hall, J., Schoenaker, D.A.J.M., Hutchinson, J., Cade, J.E., *et al.* (2018) Before the Beginning: Nutrition and Lifestyle in the Preconception Period and Its Importance for Future Health. *The Lancet*, **391**, 1830-1841.
[https://doi.org/10.1016/s0140-6736\(18\)30311-8](https://doi.org/10.1016/s0140-6736(18)30311-8)
- [23] Betran, A., Torloni, M., Zhang, J. and Gülmezoglu, A. (2015) Who Statement on Caesarean Section Rates. *BJOG: An International Journal of Obstetrics & Gynaecology*, **123**, 667-670. <https://doi.org/10.1111/1471-0528.13526>
- [24] Keag, O.E., Norman, J.E. and Stock, S.J. (2018) Long-Term Risks and Benefits Associated with Cesarean Delivery for Mother, Baby, and Subsequent Pregnancies: Systematic Review and Meta-Analysis. *PLOS Medicine*, **15**, e1002494.
<https://doi.org/10.1371/journal.pmed.1002494>
- [25] March of Dimes (2023) 2023 March of Dimes Report Card: The State of Maternal and Infant Health for American Families. March of Dimes.
<https://www.marchofdimes.org/report-card>
- [26] Frick, A.P. (2021) Advanced Maternal Age and Adverse Pregnancy Outcomes. *Best Practice & Research Clinical Obstetrics & Gynaecology*, **70**, 92-100.
<https://doi.org/10.1016/j.bpobgyn.2020.07.005>
- [27] Kenny, L.C., Lavender, T., McNamee, R., O'Neill, S.M., Mills, T. and Khashan, A.S.

- (2013) Advanced Maternal Age and Adverse Pregnancy Outcome: Evidence from a Large Contemporary Cohort. *PLOS ONE*, **8**, e56583. <https://doi.org/10.1371/journal.pone.0056583>
- [28] Šťastná, A., Fait, T., Kocourková, J. and Waldaufová, E. (2022) Does Advanced Maternal Age Comprise an Independent Risk Factor for Caesarean Section? A Population-Wide Study. *International Journal of Environmental Research and Public Health*, **20**, Article 668. <https://doi.org/10.3390/ijerph20010668>
- [29] Kozhimannil, K.B., Law, M.R. and Virnig, B.A. (2013) Cesarean Delivery Rates Vary Tenfold among US Hospitals; Reducing Variation May Address Quality and Cost Issues. *Health Affairs*, **32**, 527-535. <https://doi.org/10.1377/hlthaff.2012.1030>
- [30] Milcent, C. and Zbiri, S. (2018) Prenatal Care and Socioeconomic Status: Effect on Cesarean Delivery. *Health Economics Review*, **8**, Article No. 7. <https://doi.org/10.1186/s13561-018-0190-x>
- [31] Shah, P.S., Zao, J. and Ali, S. (2011) Maternal Marital Status and Birth Outcomes: A Systematic Review and Meta-Analyses. *Maternal and Child Health Journal*, **15**, 1097-1109. <https://doi.org/10.1007/s10995-010-0654-z>
- [32] Li, N., Liu, E., Guo, J., Pan, L., Li, B., Wang, P., et al. (2013) Maternal Prepregnancy Body Mass Index and Gestational Weight Gain on Pregnancy Outcomes. *PLOS ONE*, **8**, e82310. <https://doi.org/10.1371/journal.pone.0082310>
- [33] Braveman, P. (2023) The Black-White Disparity in Preterm Birth: Race or Racism? *The Milbank Quarterly*, **101**, 356-378. <https://doi.org/10.1111/1468-0009.12625>
- [34] Blumenshine, P., Egerter, S., Barclay, C.J., Cubbin, C. and Braveman, P.A. (2010) Socioeconomic Disparities in Adverse Birth Outcomes. *American Journal of Preventive Medicine*, **39**, 263-272. <https://doi.org/10.1016/j.amepre.2010.05.012>
- [35] Rubens, M., Ramamoorthy, V., Saxena, A., McGranaghan, P., Veleard, E. and Hernandez, A. (2022) Obstetric Outcomes during Delivery Hospitalizations among Obese Pregnant Women in the United States. *Scientific Reports*, **12**, Article No. 6862. <https://doi.org/10.1038/s41598-022-10786-9>
- [36] Zhang, Y., Lu, M., Yi, Y., Xia, L., Zhang, R., Li, C., et al. (2024) Influence of Maternal Body Mass Index on Pregnancy Complications and Outcomes: A Systematic Review and Meta-Analysis. *Frontiers in Endocrinology*, **15**, Article 1280692. <https://doi.org/10.3389/fendo.2024.1280692>
- [37] Ellis, J.A., Brown, C.M., Barger, B. and Carlson, N.S. (2019) Influence of Maternal Obesity on Labor Induction: A Systematic Review and Meta-Analysis. *Journal of Midwifery & Women's Health*, **64**, 55-67. <https://doi.org/10.1111/jmwh.12935>
- [38] Endalifer, M.L., Diress, G., Almar, H. and Endalifer, B.L. (2021) Effect of Overweight/Obesity on Caesarean Section Occurrence among Reproductive-Aged Women in Ethiopia: A Secondary Data Analysis. *BMJ Nutrition, Prevention & Health*, **4**, 111-114. <https://doi.org/10.1136/bmjnp-2020-000121>
- [39] Vinutha, B.R. and Savithri, D.R. (2025) Effect of Pre-Pregnancy Body Mass Index on Mode of Delivery: A Comprehensive Observational Study. *European Journal of Cardiovascular Medicine*, **15**, 588-594.
- [40] Xiong, C., Zhou, A., Cao, Z., Zhang, Y., Qiu, L., Yao, C., et al. (2016) Association of Pre-Pregnancy Body Mass Index, Gestational Weight Gain with Cesarean Section in Term Deliveries of China. *Scientific Reports*, **6**, Article No. 37168. <https://doi.org/10.1038/srep37168>
- [41] McHale, P., Maudsley, G., Pennington, A., Schlüter, D.K., Barr, B., Paranjothy, S., et al. (2022) Mediators of Socioeconomic Inequalities in Preterm Birth: A Systematic

Review. *BMC Public Health*, **22**, Article No. 1134.

<https://doi.org/10.1186/s12889-022-13438-9>

- [42] Siddika, N., Song, S., Margerison, C.E., Kramer, M.R. and Luo, Z. (2023) The Impact of Place-Based Contextual Social and Environmental Determinants on Preterm Birth: A Systematic Review of the Empirical Evidence. *Health & Place*, **83**, Article ID: 103082. <https://doi.org/10.1016/j.healthplace.2023.103082>
- [43] Cornish, R.P., Magnus, M.C., Urhoj, S.K., Santorelli, G., Smithers, L.G., Odd, D., *et al.* (2024) Maternal Pre-Pregnancy Body Mass Index and Risk of Preterm Birth: A Collaboration Using Large Routine Health Datasets. *BMC Medicine*, **22**, Article No. 10. <https://doi.org/10.1186/s12916-023-03230-w>